

# Non-Geometric T-duality from Higher Groupoid Bundles with Connections



Christian Saemann  
Maxwell Institute and  
School of Mathematical and Computer Sciences  
Heriot-Watt University, Edinburgh

ESI Workshop on Higher Structures, 18.8.2022

Based on joint work with Hyungrok Kim: [arXiv:2204.01783](https://arxiv.org/abs/2204.01783)

see also [Konrad's talk](#)

- (Almost) no theorems, but: **framework for T-duality**
- This story has many **loose ends**:
  - Initiated as a side project to keep my PhD student busy
  - Constructions are fine, but:
    - open mathematical questions and
    - relation to physics should be explored further
    - **Very happy about questions/comments**
- I'm not claiming to be an expert on T-duality:
  - **Very happy about comments**
- Positive: higher structures, but everything very **explicit**

- String theories on backgrounds with  $U(1)$ -isometries:  
⇒ a T-dual partner
- Low-energy limit: corresponding supergravity contains  $B$ -field:  
⇒ connective structure on a gerbe

## Geometric string background:

- A Riemannian manifold  $X$
- A principal/affine torus bundle  $\pi : P \rightarrow X$  (with connection)
- An abelian gerbe (with connection)  $\mathcal{G}$  on the total space of  $P$

Ignore dynamics, i.e. no equations of motion imposed

Geometric string background:

- A Riemannian manifold  $X$
- A principal/affine torus bundle  $\pi : P \rightarrow X$
- An abelian gerbe  $\mathcal{G}$  on the total space of  $P$

Topological T-duality follows from exactness of the Gysin sequence:

$$\dots \rightarrow H^3(X, \mathbb{Z}) \xrightarrow{\pi^*} H^3(P, \mathbb{Z}) \xrightarrow{\pi_*} H^2(X, \mathbb{Z}) \xrightarrow{F \cup} H^4(X, \mathbb{Z}) \rightarrow \dots$$

- Gerbe over  $P$ : 3-form  $H \in H^3(P, \mathbb{Z})$
- Fiber integration  $\pi_* H = \hat{F} \in H^2(X, \mathbb{Z})$  with  $F \cup \hat{F} = 0$
- $\Rightarrow$  There is  $\hat{H} \in H^3(\hat{P}, \mathbb{Z})$  with  $\pi_* \hat{H} = F$ .
- Top. T-duality:  $(F, H) = (\pi_* \hat{H}, H) \longleftrightarrow (\hat{F}, \hat{H}) = (\pi_* H, \hat{H})$

Note: possibility of topology change!

Bouwknegt, Evslin, Hannabuss, Mathai (2004)

T-correspondence:

$$\begin{array}{c}
 \mathcal{G}_C = \check{p}^* \check{\mathcal{G}} \otimes \hat{p}^* \hat{\mathcal{G}}^{-1} \cong \mathcal{I} \\
 \downarrow \\
 \check{P} \times_X \hat{P} \\
 \swarrow \check{p} \quad \searrow \hat{p} \\
 \check{\mathcal{G}} \rightarrow \check{P} \quad \hat{P} \leftarrow \hat{\mathcal{G}} \\
 \searrow \check{\pi} \quad \swarrow \hat{\pi} \\
 X
 \end{array}$$

Bunke, Rumpf, Schick (2005, 2006)

Principal 2-bundles (without connections) over  $X$ :

$$\begin{array}{c}
 \mathcal{P}_C \\
 \swarrow \check{p} \quad \searrow \hat{p} \\
 \check{\mathcal{P}} \quad \hat{\mathcal{P}}
 \end{array}$$

Nikolaus, Waldorf (2018)

## I. T-duality can lead to **non-geometric backgrounds**:

$F^3$ :  $H$  has no legs along fiber

T-duality: identity

$F^2$ :  $H$  has 1 leg along fiber

T-duality  $\rightarrow$  geometric string background

$F^1$ :  $H$  has 2 legs along fiber

T-duality  $\rightarrow$   $Q$ -space, (e.g. T-folds) locally geometric

$F^0$ :  $H$  has all legs along fiber

T-duality  $\rightarrow$   $R$ -space, non-geometric

Nikolaus/Waldorf cover  $F^2 \leftrightarrow F^2$  and  $F^2 \leftrightarrow F^1$  T-dualities

**What about the general case?**

## II. **Differential refinement** of this picture

### Why is this interesting/hard?

- I. need to use suitable **groupoids** and **augmented groupoids**
- II. connections on principal 2-bundles often require **adjustment**

- Recap: Adjusted connections on **principal 2-bundles**
- $F^k$ ,  $k \geq 2$ : **Geometric T-duality** with principal 2-bundles
- The T-duality group from **Kaluza–Klein reduction**
- $F^k$ ,  $k \geq 1$ : Groupoid bundles for **T-folds/Q-spaces**
- $F^k$ ,  $k \geq 0$ : Augmented groupoid bundles for **R-spaces**
- Explicit examples throughout: **Nilmanifolds**

Principal 2-bundles or Non-Abelian Gerbes  
with Adjusted Connections



A mathematical structure (“Bourbaki-style”) consists of

- Sets
- Structure Functions
- Structure Equations

“Categorification”:

Sets  $\rightarrow$  Categories

Structure Functions  $\rightarrow$  Structure Functors

Structure Equations  $\rightarrow$  Structure Isomorphisms

Example: Group  $\rightarrow$  2-Group

- Set  $G \rightarrow$  Category  $\mathcal{G}$
- product, identity ( $\mathbb{1} : * \rightarrow G$ ), inverse  $\rightarrow$  Functors
- $a(bc) = (ab)c \rightarrow$  Associator  $a : a \otimes (b \otimes c) \Rightarrow (a \otimes b) \otimes c$
- $\mathbb{1}a = a\mathbb{1} = a \rightarrow$  Unitors  $l_a : a \otimes \mathbb{1} \Rightarrow a$ ,  $r_a : \mathbb{1} \otimes a \Rightarrow a$
- $aa^{-1} = a^{-1}a = \mathbb{1} \rightarrow$  weak inv.  $\text{inv}(x) \otimes x \Rightarrow \mathbb{1} \leftarrow x \otimes \text{inv}(x)$

Note: Process not unique, variants: weak/strict/...

$$\mathbb{R}^{2n} \times \mathbb{Z}^{2n} \times \text{U}(1) \rightrightarrows \mathbb{R}^{2n}$$

$$\begin{array}{ccccc}
 & & (\xi, m_1, \phi_1) & & (\xi - m_1, m_2, \phi_2) \\
 & \swarrow & \leftarrow & \swarrow & \leftarrow \\
 \xi & & \xi - m_1 & & \xi - m_1 - m_2 \\
 & \swarrow & & \swarrow & \\
 & & (\xi, m_1 + m_2, \phi_1 + \phi_2) & & 
 \end{array}$$

$$\text{id}_\xi := (\xi, 0, 0), \quad (\xi, m, \phi)^{-1} := (\xi - m, -m, -\phi)$$

$$(\xi_1, m_1, \phi_1) \otimes (\xi_2, m_2, \phi_2) := (\xi_1 + \xi_2, m_1 + m_2, \phi_1 + \phi_2 - \langle \xi_1, m_2 \rangle)$$

$$\text{inv}(\xi, m, \phi) := (-\xi, -m, -\phi - \langle \xi, m \rangle)$$

This Lie 2-group corresponds to a **crossed module of Lie groups**:

$$\text{TD}_n := (\mathbb{Z}^{2n} \times \text{U}(1) \xrightarrow{\text{t}} \mathbb{R}^{2n})$$

$$\text{t}(m, \phi) := m$$

$$\xi \triangleright (m, \phi) := (m, \phi - \langle \xi, m \rangle)$$

Essentially, all definitions of principal bundles have higher version.

Here: Čech cocycle description subordinate to a cover.

- Surjective submersion  $\sigma : Y \twoheadrightarrow X$ , e.g.  $Y = \sqcup_a U_a$
- Čech groupoid:

$$\check{\mathcal{C}}(\sigma) : Y \times_X Y \rightrightarrows Y, \quad (y_1, y_2) \circ (y_2, y_3) = (y_1, y_3).$$

- Principal G-bundle:

Transition functions are functor  $g : \check{\mathcal{C}}(\sigma) \rightarrow (\mathbf{G} \rightrightarrows *)$

$$\begin{array}{ccc} Y \times_X Y & \xrightarrow{g} & \mathbf{G} \\ \Downarrow & & \Downarrow \\ Y & \xrightarrow{*} & * \end{array} \quad g(y_1, y_2)g(y_2, y_3) = g(y_1, y_3)$$

Equivalences/bundle isomorphisms: natural isomorphisms.

- 2-groupoid version yields principal 2-bundles, including gerbes.

Connections on principal 2-bundles: work a bit more...

Breen, Messing (2005), Aschieri, Cantini, Jurčo (2005)

Data obtained for 2-group  $G \times H \rightrightarrows G$  and Lie 2-algebra  $\mathfrak{g} \times \mathfrak{h} \rightrightarrows \mathfrak{g}$ :

$$h \in \Omega^0(Y^{[3]}, \mathfrak{H}) \quad \Lambda \in \Omega^1(Y^{[2]}, \mathfrak{h}) \quad B \in \Omega^2(Y, \mathfrak{h}) \quad \delta \in \Omega^2(Y^{[2]}, \mathfrak{h})$$

$$g \in \Omega^0(Y^{[2]}, \mathfrak{G}) \quad A \in \Omega^1(Y, \mathfrak{g})$$

- Note that  $\delta$  sticks out unnaturally.
- It was dropped in most later work (Baez, Schreiber, ...)
- Price to pay: **part of curvature must vanish**
- Otherwise, gauge transformations **do not compose**

Object	Principal $G$ -bundle	Principal $(H \xrightarrow{t} G)$ -bundle
Cochains	$(g_{ab})$ valued in $G$	$(g_{ab})$ valued in $G$ , $(h_{abc})$ valued in $H$
Cocycle	$g_{ab}g_{bc} = g_{ac}$	$t(h_{abc})g_{ab}g_{bc} = g_{ac}$ $h_{acd}h_{abc} = h_{abd}(g_{ab} \triangleright h_{bcd})$
Coboundary	$g_a g'_{ab} = g_{ab} g_b$	$g_a g'_{ab} = t(h_{ab})g_{ab}g_b$ $h_{ac}h_{abc} = (g_a \triangleright h'_{abc})h_{ab}(g_{ab} \triangleright h_{bc})$
gauge pot.	$A_a \in \Omega^1(U_a) \otimes \mathfrak{g}$	$A_a \in \Omega^1(U_a) \otimes \mathfrak{g}$ , $B_a \in \Omega^2(U_a) \otimes \mathfrak{h}$
Curvature	$F_a = dA_a + A_a \wedge A_a -$	$\mathcal{F}_a = dA_a + \frac{1}{2}[A_a, A_a] - t(B_a) \stackrel{!}{=} 0$ $H_a = dB_a + A_a \triangleright B_a$
Gauge trafos	$\tilde{A}_a := g_a^{-1}A_a g_a + g_a^{-1}dg_a$	$\tilde{A}_a := g_a^{-1}A_a g_a + g_a^{-1}dg_a + t(\Lambda_a)$ $\tilde{B}_a := g_a^{-1} \triangleright B_a + \tilde{A}_a \triangleright \Lambda_a + d\Lambda_a - \Lambda_a \wedge \Lambda_a$

## Remarks:

- A principal  $(1 \xrightarrow{t} G)$ -bundle is a principal  $G$ -bundle.
- A principal  $(U(1) \xrightarrow{t} 1) = BU(1)$ -bundle is an abelian gerbe.

# Why should the fake curvature(s) vanish?

$$\mathcal{F} := dA + \frac{1}{2}[A, A] + \mathfrak{t}(B) \stackrel{!}{=} 0$$

Without this condition:

- Gauge transformations **do not close**
- Cocycles **do not glue together**
- Higher parallel transport **is not reparameterization invariant**
- 6d Self-duality equation  $H = \star H$  **is not gauge-covariant:**

$$H \rightarrow \tilde{H} = g \triangleright H - \mathcal{F} \triangleright \Lambda$$

With this condition:

- Principal  $(1 \xrightarrow{\mathfrak{t}} \mathbb{G})$ -bundle is **flat** principal  $\mathbb{G}$ -bundle.
- Higher connections are **locally abelian!**

Gastel (2019), CS, Schmidt (2020)

Many (not all!) higher gauge groups come with

Adjustment of higher group  $\mathcal{G}$ :

CS, Schmidt (2020), Rist, CS, Wolf (2022)

- Additional map  $\kappa : \mathcal{G} \times \text{Lie}(\mathcal{G}) \rightarrow \text{Lie}(\mathcal{G})$  + condition
- Necessary for consistent definition of invariant polynomials.
- From Alternator ( $\Rightarrow EL_\infty$ -algebras, Borsten, Kim, CS (2021))

For connections on principal  $\mathcal{G}$ -bundles:

- specifies  $\delta \in \Omega^2(Y^{[2]}, \mathfrak{h})$  in terms of  $A$  and  $F$
- Adjustment of curvature/cocycle/coboundary relations
- Can drop fake flatness condition

Archetypal example: **string Lie 2-algebra**

$$\mathbf{string}(n) = \mathbb{R}[1] \rightarrow \mathbf{spin}(n)$$

$$\mu_2(x_1, x_2) = [x_1, x_2], \quad \mu_3(x_1, x_2, x_3) = (x_1, [x_2, x_3])$$

**Gauge potentials:**

$$(A, B) \in \Omega^1(U) \otimes \mathbf{spin}(n) \oplus \Omega^2(U)$$

**Curvatures:**


$$\begin{aligned} F &:= dA + \frac{1}{2}[A, A] \\ H &:= dB - \frac{1}{3!}\mu_3(A, A, A) + (A, F) \\ &= dB + \underbrace{(A, dA) + \frac{1}{3}(A, [A, A])}_{\mathbf{cs}(A)} \end{aligned}$$

**Bianchi identities:**

$$dF + [A, F] = 0, \quad dH - (F, F) = 0$$



## Geometric T-duality



- Nikolaus/Waldorf: Topological part:**
  - $\check{\mathcal{P}}$  and  $\hat{\mathcal{P}}$  are principal  $\mathrm{TB}_n^{\mathrm{F}2}$ -bundles
  - $\mathcal{P}_C$  is a principal  $\mathrm{TD}_n$ -bundle
  - $\hat{\rho}$  is a projection induced by strict morphism  $\hat{\phi} : \mathrm{TD}_n \rightarrow \mathrm{TB}_n^{\mathrm{F}2}$
  - $\check{\rho}$  induced by  $\check{\phi} = \hat{\phi} \circ \phi_{\mathrm{flip}}$ , flip morphism  $\phi_{\mathrm{flip}} : \mathrm{TD}_n \rightarrow \mathrm{TD}_n$
- Differential refinement:** (i.e.  $B$ -field+metric) **Kim, CS (2022)**
  - $\mathrm{TB}_n^{\mathrm{F}2}$  does not come with adjustment, but
  - $\mathrm{TD}_n$  comes with very natural adjustment map
  - Have **topological** and **full connection data** on  $\mathcal{P}_C$
  - Can reconstruct gerbe and bundle data on  $\check{\mathcal{P}}$  and  $\hat{\mathcal{P}}$
- Reproduces Buscher rules **Waldorf (2022)**
- Generalization to **affine torus bundles**: use  $\mathrm{GL}(n, \mathbb{Z}) \ltimes \mathrm{TD}_n$

Geometry of string background  $\check{\mathcal{G}}_\ell \rightarrow N_k$ :

- **Principal circle bundle** over  $T^2$  with  $c_1 = k$
- Subordinate to  $\mathbb{R}^2 \rightarrow T^2$  and with  $U(1) \cong \mathbb{R}/\mathbb{Z}$   
 $(x, y, z) \sim (x, y + 1, z) \sim (x, y, z + 1) \sim (x + 1, y, z - ky)$
- **Local connection form**:  $A(x, y) = kx \, dy \in \Omega^1(\mathbb{R}^2)$
- **Kaluza–Klein metric**:  $g(x, y, z) = dx^2 + dy^2 + (dz + kx \, dy)^2$
- Gerbes on  $N_k$  characterized by element of  $H^3(N_k, \mathbb{Z}) \cong \mathbb{Z}$

T-duality:

$$(\check{\mathcal{G}}_\ell \rightarrow N_k) \longleftrightarrow (\hat{\mathcal{G}}_k \rightarrow N_\ell)$$

Kim, CS (2022)

$$\begin{array}{ccc}
 & \mathcal{P}_C & \\
 \check{\mathcal{P}} & \xleftarrow{\check{p}} & \\
 & & \hat{\mathcal{P}} \\
 & \xrightarrow{\hat{p}} & 
 \end{array}$$

Lie 2-group:

$$\text{TD}_1 := (\mathbb{Z}^2 \times \text{U}(1) \xrightarrow{t} \mathbb{R}^2)$$

Topological cocycle data:

$$g = \begin{pmatrix} \hat{\xi} \\ \check{\xi} \end{pmatrix}, \quad \begin{aligned} \hat{\xi}(x, y; x', y') &= \ell(x' - x)y, \\ \check{\xi}(x, y; x', y') &= k(x' - x)y, \end{aligned}$$

$$h = \begin{pmatrix} \hat{m} \\ \check{m} \\ \phi \end{pmatrix}, \quad \begin{aligned} \hat{m}(x, y; x', y'; x'', y'') &= -\ell(x'' - x')(y' - y) \\ \check{m}(x, y; x', y'; x'', y'') &= -k(x'' - x')(y' - y) \\ \phi &= \frac{1}{2}k\ell(y'(xx'' - xx' - x'x'') - (x'' - x')(y'^2 - y^2)x) \end{aligned}$$

Cocycle data of differential refinement:

$$A = \begin{pmatrix} \check{A} \\ \hat{A} \end{pmatrix} = \begin{pmatrix} kx \, dy \\ \ell x \, dy \end{pmatrix}, \quad B = 0, \quad \Lambda = \frac{1}{2}k\ell(xx' \, dy + (xy + x'y' + y^2(x' - x)) \, dx)$$

Can **reconstruct** both string backgrounds fully.

## The T-duality group from Kaluza–Klein Reduction

## Observation:

T-duality is intimately linked to Kaluza–Klein reduction:

- Gysin sequence contains **fiber integration**
- Metric on total space given by **Kaluza–Klein metric**
- Literature: e.g. **Berman (2019)**, **Alfonsi (2019)**, ...
  
- Geometric objects from maps into **classifying spaces  $\mathcal{C}$** .
- Note: **currying**  $C^0(X \times T^n, \mathcal{C}) \cong C^0(X, C^0(T^n, \mathcal{C}))$
- Non-trivial fibrations: **cyclic torus space**:  $C^0(T^n, \mathcal{C}) // \mathrm{U}(1)^n$   
cf. **Fiorenza, Sati, Schreiber (2016a, 2016b)**

$TD_1$  from KK-reduction of gerbe on circle bundle

- Gerbe:  $C^0(P, \mathcal{C})$  with  $\mathcal{C} = BBU(1)$
- Cyclic loop space:  $LBBU(1)//U(1) \cong B(BU(1) \times U(1) \times U(1))$
- Replace  $U(1)$  with  $\mathbb{Z} \rightarrow \mathbb{R}$ :  $TD_1 := (U(1) \times \mathbb{Z}^2 \xrightarrow{t} \mathbb{R}^2)$

$TD_2$  from KK-reduction of principal  $TD_1$ -bundle on circle bundle

- Principal 2-bundle:  $C^0(P, \mathcal{C})$  with  $\mathcal{C} = BTD_1$
- Note:  $LBU(1)//U(1) \cong BU(1) \times U(1) \times U(1)$
- Replace  $U(1)$  with  $\mathbb{Z} \rightarrow \mathbb{R}$ :  $TD_2 := (U(1) \times \mathbb{Z}^4 \xrightarrow{t} \mathbb{R}^4)$

Iterate:  $TD_n := (U(1) \times \mathbb{Z}^{2n} \xrightarrow{t} \mathbb{R}^{2n})$

Abstract nonsense:

- Natural definition of **morphism of 2-groups**
- **Automorphisms** of 2-group form naturally a 2-group
- **2-group action**  $\mathcal{G} \curvearrowright \mathcal{H}$ : morphism  $\mathcal{G} \rightarrow \text{Aut}(\mathcal{H})$

Automorphisms of the 2-group  $\mathbf{TD}_n$ :

- Can be computed to be weak (unital) Lie 2-group

$$\mathcal{GO}(n, n; \mathbb{Z}) := \left( \text{GO}(n, n; \mathbb{Z}) \times \mathbb{Z}^{2n} \rightrightarrows \text{GO}(n, n; \mathbb{Z}) \right)$$

see also Waldorf (2022)

- While  $\text{GO}(n, n; \mathbb{Z})$  **does not** act on  $\mathbf{TD}_n$ ,  $\mathcal{GO}(n, n; \mathbb{Z})$  does.
- **Recover T-duality group** for affine torus bundles
- Explicit: **geometric subgroup**,  $B$ - and  $\beta$ -trafos, T-dualities as endo-2-functors on  $\mathbf{TD}_n$
- $\Rightarrow$  arrange everything based on  $\mathcal{GO}(n, n; \mathbb{Z})$



## Groupoid bundles for T-folds

Recall:  $\mathbf{TD}_2$  from KK-reduction of principal  $\mathbf{TD}_1$ -bundle

- $C^0(P, \mathcal{C})$  with  $\mathcal{C} = \mathbf{BTD}_1 \cong \mathbf{BBU}(1) \times \mathbf{BU}(1) \times \mathbf{BU}(1)$
- $\mathbf{LBBU}(1)//\mathbf{U}(1) \cong \mathbf{BBU}(1) \times \mathbf{BU}(1) \times \mathbf{BU}(1)$
- $\mathbf{LBU}(1)//\mathbf{U}(1) \cong \mathbf{BU}(1) \times \mathbf{U}(1) \times \mathbf{U}(1)$
- $\mathbf{BTD}_2 \cong \mathbf{BBU}(1) \times \mathbf{BU}(1)^{\times 4}$
- Replace  $\mathbf{U}(1)$  with  $\mathbb{Z} \rightarrow \mathbb{R}$ :  $\mathbf{TD}_2 := (\mathbf{U}(1) \times \mathbb{Z}^4 \xrightarrow{t} \mathbb{R}^4)$
- Here, we dropped parts, we actually get a 2-groupoid:  
 $\mathcal{TD}_2 \cong \mathbf{BBU}(1) \times \mathbf{BU}(1)^{\times 4} \times \mathbf{U}(1)^{\times 4}$
- Clear that  $g, B$  dim reduced on  $T^2$  yield four scalar modes.

For T-folds: at least **two** T-duality directions  $\Rightarrow$  2-groupoid!

- Two T-dualities yield **scalars** from metric and 2-form.
- Scalars live on the **Narain moduli space** for affine torus bundles:

$$\begin{aligned} GM_n &= \mathrm{GO}(n, n; \mathbb{Z}) \setminus \mathrm{O}(n, n; \mathbb{R}) / (\mathrm{O}(n; \mathbb{R}) \times \mathrm{O}(n; \mathbb{R})) \\ &=: \mathrm{GO}(n, n; \mathbb{Z}) \setminus Q_n \end{aligned}$$

- Note:  $Q_n \cong \mathbb{R}^{n^2}$  is a nice space
- Resolve into **action groupoid**:

$$\mathrm{GO}(n, n; \mathbb{Z}) \ltimes Q_n \rightrightarrows Q_n$$

- Extend to  $\mathcal{GO}(n, n; \mathbb{Z})$ -action ( $\mathcal{GO}(n, n; \mathbb{Z}) \cong \mathrm{Aut}(\mathrm{TD}_n)$ )
- Place  $\mathrm{TD}_n$ -fiber over every point in  $Q_n$
- Include action of  $\mathcal{GO}(n, n; \mathbb{Z})$  on  $\mathrm{TD}_n$
- The result is the 2-groupoid  $\mathcal{TD}_n$

Recall: functorial description of (higher) principal bundles:

- Manifold  $X$
- Cover/surjective submersion  $\sigma : Y \rightarrow X$
- Cech groupoid  $\check{\mathcal{C}}(\sigma) := (Y \times_X Y \rightrightarrows Y)$
- Top. principal  $G$ -bundle: functor  $\check{\mathcal{C}}(\sigma) \rightarrow \mathbf{BG}$

For  $\mathcal{TD}_n$ -bundle:

- Replace (higher) group  $\mathbf{BG}$  by Lie 2-groupoid  $\mathcal{TD}_n$
- For ordinary groupoids: e.g. gauged sigma models

A non-geometric T-duality is simply a  $\mathcal{TD}_n$ -bundle.

## Remarks:

- The T-duality group  $\mathcal{GO}(n, n; \mathbb{Z}) \supset \text{GO}(n, n; \mathbb{Z})$  is **gauged!**
- Explicitly visible:  $\text{GO}(n, n; \mathbb{Z})$ -gluing of local data
- **Matches topological discussion** in **Nikolaus, Waldorf (2018)**
- Differential refinement imposes **restriction on top. cocycles**
- This describes all T-dualities between pairs of **T-folds**
- Concrete conditions for **“half-geometric”** T-dualities
- **Concrete cocycles** of the T-fold in the nilmanifold example

To describe  $Q$ -spaces/T-folds:  
(can) use **higher** instead of **noncommutative geometry**.

Consider again the nilmanifold example, this time  $X = S^1$ .

- Gauge groupoid  $\mathcal{T}\mathcal{D}_2$
- General cocycle data:

$$(g, z, \xi, m, \phi, q) \in C^\infty(Y^{[3]}, \mathrm{GO}(2, 2; \mathbb{Z}) \times \mathbb{Z}^4 \times \mathbb{R}^4 \times \mathbb{Z}^4 \times \mathrm{U}(1) \times Q_2)$$

$$(g, \xi, q) \in C^\infty(Y^{[2]}, \mathrm{GO}(2, 2; \mathbb{Z}) \times \mathbb{R}^4 \times Q_2)$$

$$q \in C^\infty(Y, Q_2)$$

- Topology: all data over  $Y^{[3]}$  are **trivial**.
- Topology: no  $T^m$ -bundles over  $S^1$ :  $\xi$  is **trivial**
- Remaining:  $q : Y \rightarrow Q_2 \cong \mathbb{R}^4$ ,  $g : Y^{[2]} \rightarrow \mathrm{GO}(2, 2; \mathbb{Z})$  s.t.:

$$q(y_1) = g(y_1, y_2)q(y_2), \quad g(y_1, y_2)g(y_2, y_3) = g(y_1, y_3)$$

- $\mathbb{R}^4$ : scalar modes  $g_{yy}, g_{yz}, g_{zz}, B_{yz}$
- **Well-known T-fold** is the special case where

$$g_{x+1,x} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & \ell & 1 & 0 \\ -\ell & 0 & 0 & 1 \end{pmatrix}$$

## Augmented groupoid bundles for $\mathbb{R}$ -spaces

- T-folds/ $Q$ -spaces relatively harmless, as **locally geometric**
- $R$ -spaces are not even locally geometric
- But perhaps **higher description** still works?

Note:

- **One** T-duality direction:  $B$ -field  $\rightarrow$  2-, 1-forms  
 $\Rightarrow$  Lie 2-group  $TD_n$ -bundles with connection
- **Two** T-duality directions:  $B$ -field  $\rightarrow$  2-, 1-, 0-forms  
 $\Rightarrow$  Lie 2-groupoid  $\mathcal{T}\mathcal{D}_n$ -bundles with connection
- **Three** T-duality directions:  $B$ -field  $\rightarrow$  2-, 1-, 0-, “(-1)-forms”  
(Note: (-1)-forms have global “curvature” 0-forms)  
 $\Rightarrow$  **Augmented** Lie 2-groupoid  $\mathcal{T}\mathcal{D}_n^{\text{aug}}$ -bundles with connection



Need to switch to **simplicial picture**:

- (Higher) groupoids are **Kan simplicial manifolds**
- Higher groupoid 1-morphisms are **simplicial maps**
- Higher groupoid 2-morphisms are **simplicial homotopies**
- “**quasi-groupoids**” or “ **$(\infty, 1)$ -groupoids**”

**Augmented  $\mathcal{G}$ -groupoid bundles** subordinate to  $\sigma : Y \twoheadrightarrow X$ :

$$\begin{array}{ccc}
 Y \times_X Y \times_X Y & \xrightarrow{g_2} & \mathcal{G}_2 \\
 \Downarrow & & \Downarrow \\
 Y \times_X Y & \xrightarrow{g_1} & \mathcal{G}_1 \\
 \Downarrow & & \Downarrow \\
 Y & \xrightarrow{g_0} & \mathcal{G}_0 \\
 \downarrow \sigma & & \downarrow \\
 X & \xrightarrow{g_{-1}} & \mathcal{G}_{-1}
 \end{array}$$

Construction of  $\mathcal{I}\mathcal{D}_n^{\text{aug}}$ :

- Augmentation by suitable space of  $R$ -fluxes
- Determined by finite version of **tensor hierarchy**
- Finite **embedding tensor**  $\mathbb{R}^{2n} \rightarrow \text{GO}(n, n; \mathbb{Z}) \subset \mathcal{G}\mathcal{O}(n, n; \mathbb{Z})$
- plus some standard consistency conditions
- Beyond this, augmentation **fairly trivial**

Remarks on T-duality with  $\mathcal{I}\mathcal{D}_n^{\text{aug}}$ -bundles:

- **Explicit examples**, e.g. from nilmanifolds
- Yields **consistency conditions** between  $Q$ - and  $R$ -fluxes
- **All previously discussed** cases included
- **All previously discussed** also for affine  $U(1)$ -bundles

To describe  $R$ -spaces:  
(can) use **higher** instead of **nonassociative geometry**.

What has been done:

- Top. T-duality can be described using **principal 2-bundles**
- Differential refinement with **adjusted curvatures**
- Explicit description of geometric T-duality with **nilmanifolds**
- T-duality group is really a 2-group derived from **KK-reduction**
- Extended to  **$Q$ -spaces** or **T-folds**
- Extended to  **$R$ -spaces**

Future work:

- Link some mathematical results to **physical expectations**
- Link to **pre- $NQ$ -manifold pictures**, DFT, and similar
- Non-abelian T-duality?
- **U-duality**

Thank You!